Raa

#### **PCT**

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:
A23L 1/308, 1/30, 1/305

A1

(11) International Publication Number: WO 99/53777

(43) International Publication Date: 28 October 1999 (28.10.99)

(21) International Application Number: PCT/EP99/02668

(22) International Filing Date: 20 April 1999 (20.04.99)

(71) Applicant (for all designated States except US): NOVARTIS NUTRITION AG [CH/CH]; Monbijoustrasse 118, CH-3001 Bern (CH).

22 April 1998 (22.04.98)

(72) Inventors; and
(75) Inventors/Applicants (for US only): LUSTENBERGER, Alice
[CH/CH]; Murtenstrasse 226, CH-3207 Bern (CH). CLAY-TON, Diane, Elizabeth [GB/CH]; Au Charmin, CH-1648

(74) Agent: SMOLDERS, Walter, Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### **Published**

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: NUTRITIONAL SUPPLEMENT FOR PREGNANT AND LACTATING WOMEN

#### (57) Abstract

(30) Priority Data:

9808579.8

Hauteville (CH).

A nutritional composition for pregnant and lactating women comprising: a non-milk-derived protein source, a dietary fibre and/or fibre-like substance, a source of polyunsaturated fatty acids, minerals and vitamins recommended for pregnant and lactating women, characterized in that the composition is lactose free and contains less than 2 % of milk or milk-derived products by weight of dry components of the composition.

### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
M	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΛZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL.	Tsrael	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	υz	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Câte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

#### Nutritional Supplement for Pregnant and Lactating Women

This invention provides a nutritional composition for pregnant and lactating women and food products or dietary supplements comprising said composition.

During pregnancy and lactation a woman's diet has to provide sufficient nutrients both for the maintenance of her own body and the growth of the foetus in her womb and/or the nursing baby. The diet during such periods should be very well balanced and cover the needs of all essential nutrients. When the extra-needs of essential nutrients are compared with the extra-needs of energy during the pregnancy and lactation, it can be seen that the needs of essential nutrients increase by 50% or more whereas the energy needs increase only 15-25%. Therefore, if the diet of the pregnant and lactating women does not change in quality the extra-needs of essential nutrients might not be covered. Administering a specifically tailored "nutrient dense" nutritional composition helps ensure that the extraneeds of essential nutrients during pregnancy and lactation are covered.

Constipation is common in pregnancy. This is thought to be related to the increase in circulating progesterone and possibly decreased levels of motilin. Dietary fibre has a major effect on colonic function and is a safe means to relieve pregnant women of their constipation.

The important role of essential fatty acids in neural development of the foetus and young infant is becoming clearer and because disorders of brain development can be permanent, proper provision should be made to protect the long chain polyunsaturated fatty acid status of both term and preterm infants to ensure optimum conditions for the development of membrane-rich systems such as the brain, nervous, and vascular system. Adequate levels of long chain polyunsaturated fatty acids in the maternal diet even shortly before and during pregancy and during lactation are, therefore, important to avoid nutritional inadequacies of the foetus and nursing baby.

Calcium is among the nutrients the recommended levels of which is increased proportionately more than that of energy and the other nutrients during pregnancy and lactation and taking extra milk or dairy products will normally provide additional calcium.

-2-

This recommendation cannot be followed by all women. In particular lactose intolerant individuals need an alternative calcium source which contains no lactose. Lactose-intolerance is one of the major reasons for milk intolerance. Furthermore, it is known that exposure of the foetus via the placenta to cow's milk can result in primary sensitisation to cow's milk allergens, leading to positive proliferative responses when the infant is exposed to cow's milk after birth.

It is an object of the present invention to provide a nutritional composition which supplies the extra-needs of essential nutrients for pregnant and lactating women and which is lactose-free and which is essentially free of milk products and is, therefore, suitable for women who are lactose intolerant and women expecting a baby with an increased risk of developing cow's milk allergy because of i.a. a family history of allergies.

The present invention, therefore, provides a nutritional composition for pregnant and lactating women comprising:

- a non-milk-derived protein source,
- a dietary fibre and/or fibre-like substance,
- a source of polyunsaturated fatty acids,
- minerals and vitamins recommended for pregnant and lactating women, characterized in that the composition is lactose-free and contains less than 2 % of milk or milk-derived products by weight of dry components of the composition.

Preferably the composition according to the invention contains less than 1.5 weight % (more preferred less than 1 weight%) by weight of dry components of the composition of milk or milk-derived products such as caseinates, particularly preferred the composition is milk-free, i.e. it contains no detectable levels of milk or milk-derived products. Lactose-free as used herein means that the composition contains no detectable lactose.

Preferably the composition according to the invention comprises the following minerals and vitamins recommended for pregnant and lactating women: calcium, magnesium, iron, copper, zinc, iodine, vitamin A, vitamin E, vitamin B<sub>1</sub>, vitamin B<sub>2</sub>, vitamin B<sub>6</sub>, vitamin B<sub>12</sub>, vitamin C, folic acid, niacin.

- 3 -

PCT/EP99/02668

The composition of the invention preferably comprises the following daily amounts of essential nutrients:

- 1.0 mg (preferably 1.0 g) to 50 g of a non-milk-derived protein source, preferably 8 to 15 g, particularly preferred 10 g,
- 1.0 mg to 30 g of dietary fibre and/or fibre-like substance, preferably 4 to 6 g, particularly preferred 5 g,

of polyunsaturated fatty acids

- 0.01 μg (preferably 0.1 μg) to 2.0 g of long-chain polyunsaturated fatty acids, preferably 0.1 to 0.4 g, particularly preferred 0.25 g and/or
- 0.5 g to 12 g of alpha-linolenic acid, preferably 0.5 to 2.0 g, particularly preferred 1.0 g and/ or
- 1 g to 30 g of linoleic acid, preferably 2.0 to 5.0 g, particularly preferred 3.0 g, 0.1 mg to 3.6 g calcium, preferably 320 to 530 mg, particularly preferred 400 mg, 0.1 mg to 1.1 g magnesium, preferably 60 to 90 mg, particularly preferred 75 mg, 0.01 mg to 90 mg iron, preferably 8 to 12 mg, particularly preferred 10 mg, 0.01 mg to 60 mg zinc, preferably 4 to 6 mg, particularly preferred 5 mg, 0.01 mg to 3 mg copper, preferably 0.1 to 0.4 mg, particularly preferred 0.25 mg, 0.01 mg to 0.6 mg iodine, preferably 24 to 36 μg, particularly preferred 30 μg, 0.01 mg to 4.0 mg vitamin A, preferably 80 to 120 μg, particularly preferred 100 μg, 0.01 mg to 40 mg vitamin E, preferably 2.4 to 3.6 mg, particularly preferred 3 mg, 0.01 mg to 5 mg vitamin B<sub>1</sub>, preferably 0.16 to 0.24 mg, particularly preferred 0.2 mg, 0.01 mg to 6 mg vitamin B<sub>2</sub>, preferably 0.32 to 0.48 mg, particularly preferred 0.4 mg, 0.01 mg to 7 mg vitamin B<sub>6</sub>, preferably 0.24 to 0.36 mg, particularly preferred 0.3 mg, 0.01 μg to 8 μg vitamin B<sub>12</sub>, preferably 0.4 μg to 0.6 μg, particularly preferred 0.5 μg,
- 0.01 mg to 300 mg vitamin C, preferably 20 to 30 mg, particularly preferred 25 mg,
- 1.0  $\mu g$  to 1.2 mg folic acid, preferably 160 to 240  $\mu g$ , particularly preferred 200  $\mu g$ , and
- 0.01 mg to 60 mg niacin, preferably 1.6 to 2.4 mg, particularly preferred 2 mg.

The following sources are suitable as non-milk-derived protein source: egg protein, soy protein (particularly soy protein low in isoflavones), soy isolate, pea protein, pea hydrolysate, rice protein, rice hydrolysate, soy hydrolysate, wheat hydrolysate, potato

- 4 -

PCT/EP99/02668

hydrolysate, rapeseed protein, lupin protein, wheat protein and/or wheat isolate. Preferably the non-milk-derived protein source is soy protein.

Suitable dietary fibres include insoluble and soluble fibres and mixtures thereof. Suitable insoluble fibres include one or more members selected from the following group: cellulose, wheat fibre, soy fibre, pea fibre, oat fibre and beet fibre. Suitable soluble fibres include one or more members selected from the following group: pectin, xanthan gum, hydrolysed guar gum and gum arabic. Suitable substances with fibre-like activity are inulin and oligosaccharides such as hydrolysed inulin, soyoligosaccharides, galactooligosaccharides, fructo-oligosaccharides and xylo-oligosaccharides. Oligosaccharides are prebiotic substances with fibre-like physiological activity, i.e. they have mild laxative effect. Preferably the composition comprises oligosaccharides and in particular fructooligosaccharides and/or xylo-oligosaccharides as fibre-like substances.

For mixtures of soluble fibers and/or fibre-like substances with insoluble fibers the weight ratios are as follows:

a) soluble and/or fiber-like substance to insoluble fiber can vary from 9:1 to 2:3, a preferrred weight ratio is about 2-4 part soluble fibers and/or fibre-like substance to about 1 part insoluble fibers.

Suitable polyunsaturated fatty acids (PUFAs) include the long-chain PUFAs (from C20 upwards) such as docosahexaenoic acid (DHA), docosapentaenoic acid (DPA), eicosapentaenoic acid (EPA), arachidonic acid (AA), or their precursors alpha-linolenic acid, linoleic acid and/or docosatetraenoic acid (DTA= Adrenic), whereby the long-chain PUFAs DHA and/or EPA are preferred. Preferred sources of long chain polyunsaturated fatty acids are fish oil which typically contains 15 to 25% DHA and 5 to 15% EPA as well as sources from microalgae which typically contain 40% DHA. If the long-chain polyunsaturated fatty acid employed is of the precursor-type which still has to be metabolized to the active form in the body such as alpha-linolenic acid or linoleic acid, then the daily amount preferably lies in the range of 0.5 g to 12 g or 1 g to 30 g respectively. If the long-chain polyunsaturated fatty acid is already in the active form such as DHA or EPA then the daily amount preferably lies in the range of 0.01 µg (preferably 1.0 µg) to 2.0 g of long-chain poly-unsaturated fatty acid.

- 5 -

Suitable calcium forms are e.g. calcium phosphate, calcium citrate, calcium carbonate or calcium amino acid chelate.

Suitable magnesium forms are e.g. magnesium oxide or magnesium sulfate.

Suitable iron forms are e.g. iron pyrophosphate, iron polymaltose, iron sulphate or iron chelate.

Suitable zinc forms are e.g. zinc sulphate or zinc lactate.

Suitable copper forms are e.g. copper gluconate, copper sulphate or copper oxide.

Suitable iodine forms are e.g. potassium iodide or potassium iodate.

Preferably the nutritional composition further comprises one or more of the following minerals and vitamins: potassium, selenium, manganese, vitamin K, biotin, pantothenic acid, choline, vitamin D3 and beta-carotene. Typical daily amounts of these minerals and vitamins are 0.1 mg to 5 g of potassium, 0.1  $\mu$ g to 200  $\mu$ g of selenium, 0.1  $\mu$ g to 500  $\mu$ g manganese, 0.1  $\mu$ g to 200  $\mu$ g vitamin K, 0.1  $\mu$ g to 150  $\mu$ g biotin, 0.1  $\mu$ g to 5 mg panthotenic acid, 0.1 mg to 1 g choline, 0.1  $\mu$ g to 30  $\mu$ g of vitamin D3 and 0.1 mg to 30 mg of beta-carotene.

Suitable potassium forms are e.g. potassium citrate or potassium phosphate.

A suitable selenium form is e.g. selenium yeast.

A suitable manganese form is e.g. manganese phosphate.

The nutritional composition preferably further comprises a carbohydrate source. Typically daily amounts of 20 g to 70 g of a carbohydrate source are provided.

Suitable carbohydrate sources include maltodextrins, malt extract, cereals, fruit flakes, invert sugar, sucrose, fructose, dextrose and glucose, whereby maltodextrin and cereals are preferred.

PCT/EP99/02668

- 6 -

The nutritional composition may be in the form of a soup or drink such as a powdered cold or hot water instant drink or soup or a ready-to-consume liquid form, in the form of a bar such as. a cereal or fruit bar, in form of a salted or sweet biscuit or in the form of a meal replacement.

Preferably the composition of the invention provides the daily amounts of essential nutrients in a single dose unit.

The nutritional compositions of the present invention can be formulated into food products or dietary supplements.

Therefore the invention also relates to food products comprising the nutritional composition of the invention. The term "food product" is intended to cover the whole variety of foods and beverages, including but not limited to powdered instant drinks, baked products such as bread, crisp bread sandwiches, biscuits and cakes, dairy substitute foods, desserts, confectionary products, spreads, cereal and/or fruit bars, breakfast cereals, juices and the like. Preferred food formats are powdered instant drinks and cereal bars.

Preferred as food products are those which are likely to be classified as "functional foods", i.e. foods that are similar in appearance to conventional foods and are intended to be consumed as part of a normal diet or as a supplement, but have been nutritionally tailored to perform physiological roles beyond the provision of simple nutrient requirements.

The invention also foresees dietary supplements comprising the composition of the invention. Typically, the latter are intended for ingestion in pill, capsule, tablet, chewable candies, tabs or liquid form.

Also provided is the use of the composition of the invention in the manufacture of a food product or dietary supplement for pregnant and lactating women.

The invention further provides a method for supplementing the extra-needs of essential nutrients of pregnant and lactating women and for treating and/or preventing constipation comprising administering to said pregnant and lactating women effective amounts of

-7-

a nutritional composition comprising:

- a non-milk-derived protein source,
- a dietary fibre and/or fibre-like substance,
- a source of polyunsaturated fatty acids,

minerals and vitamins recommended for pregnant and lactating women, whereby the composition is lactose-free and contains less than 2 % of milk or milk-derived products by weight of dry components of the composition.

The invention will be further illustrated by the following examples.

#### Example 1

A powdered cold water instant drink, to be reconstituted by 35g powder in 150 ml cold water which has following profile: Recommended daily amount 35g

soya isolate	34.0%
sucrose	18.0%
fructo-oligosaccharides	15.6%
encapsulated fish oil, BASF	9.6%
maltodextrin DE20	8.6%
cocoa powder	7.0%
mineral mix	4.0%
chocolate flavour	1.5%
MCT oil	1.0%
vitamin mix	0.4%
vanilla	0.3%

All ingredients are passed through a sieving machine with sieve (mesh size 1.45mm). The MCT oil is added to the maltodextrin and mixed for 10 minutes in a Carpentier mixer. Then the remaining ingredients are added to the maltodextrin blend and further mixed for 20 minutes.

With the following mineral mix (MM) composition:

maltodextrin DE20

54.6865 %

-8-

calcium-tri-phosphate	30.0%
magnesium oxide, light	10.2%
iron pyrophosphate	3.8%
zinc sulfate	1.15%
copper gluconate	0.16%
potassium iodide	0.0035%

All ingredients of the mineral mix are passed through a sieving machine with sieve (mash size 1.45mm), then mixed in a Ruberg mixer for 15 minutes.

## With the following vitamin mix (VM)

maltodextrin DE 20	44.203%
Vitamin C	30%
beta carotene 10%	12.7%
Vitamin E acetate 50%	7.4%
Niacinamide	1.85%
Vitamin D3	1.75%
Vitamin A acetate 325	0.95%
Vitamin B₂	0.4%
Vitamin B <sub>6</sub> HCI	0.29%
Folic acid	0.21%
Vitamin B <sub>1</sub> nitrate	0.2%
Vitamin B <sub>12</sub> 1%	0.047%

All ingredients of the vitamin mix are passed through a sieving machine with sieve (mash size 1.45mm), then mixed in a Ruberg mixer for 15 minutes.

### Example 2

A cereal bar with the following profile: Recommended daily amount 40 g

glucose syrup	25.0 %
invert sugar	8.0 %
oat flakes	16.29%
crisp rice	10.0 %

WO 99/53777	PCT/EP99/02668
- 9 -	
•	
soya isolate	12.0 %
fructo-oligosaccharides	10.0 %
banana flakes	7.5 %
sunflower oil	7.5 %
MM (as in Example 1)	1.2 %
VM (as in Example 1)	0.5 %
source of microalgae oil DHASCO, Martek Biosciences Corporation	1.5 %
soya lecithin	0.5 %
antioxidant vit E	0.01 %

The glucose syrup and inverted sugar are heated up to 110°C. The sunflower oil is first warmed separately to 50°C in a tank, then the soya lecithin and antioxidant are added to the sunflower oil and mixed until homogenous. The oil blend is added to the syrup mix and mixed in a dough mixer for 10 minutes. The microalgal oil is added and further mixed for 5 minutes. All the remaining ingredients are added and further mixed for 10 minutes. Then the mass is rolled out, cooled down to 20°C and formed into bars.

#### Example 3

A powdered cold water/juice instant drink, to be reconstituted by 32g powder in 150-200 ml cold water or fruit juice which has following profile: Recommended daily amount 32g

#### Composition:

pea protein	19.0%
sucrose	32.36%
inulin	13.8%
encapsulated fish oil, BASF	8.6%
maltodextrin DE20	9.6%
banana powder	5.0%
oat fiber	4.0%
calcium lactate, soluble	3.32
banana flavour	1.6%
Mg-glycerophosphate	0.85%
Ca-H-phosphate	0.5%

WO 99/53777	PCT/EP99/02668
-------------	----------------

- 10 -

mineral mix	0.46%
choline-H-tartrate	0.38%
vitamin mix	0.26%
xanthan gum	0.25%
beta-carotene	0.02%

## With the following mineral mix composition:

maltodextrin DE20	48.664 %
iron-III-pyrophosphate	39.46%
selenium yeast 1000	6.5%
zinc sulphate	5.35%
potassium iodide	0.026%

All ingredients of the mineral mix are passed through a sieving machine with sieve (mesh size 1.45mm), then mixed in a Ruberg mixer for 15 minutes.

## With the following vitamin mix

maltodextrin DE 20	13.606%
Vitamin C	65%
Vitamin K <sub>1</sub> powder 5%	0.15%
Vitamin E acetate 50%	8.3%
Niacinamide	2.7%
Vitamin D3	2.8%
Vitamin A acetate 325	4.8%
Vitamin B₂	0.42%
Vitamin B <sub>6</sub> HCl	1.5%
Folic acid	0.001%
Vitamin B <sub>1</sub> nitrate	0.65%
Vitamin B <sub>12</sub> 1%	0.073%

All ingredients of the vitamin mix are passed through a sieving machine with sieve (mash size 1.45mm), then mixed in a Ruberg mixer for 15 minutes.

PCT/EP99/02668

- 11 -

### Example 4

A liquid product with the following profile: Recommended daily portion 200 ml

water	80.09%
sucrose	7.5%
soya isolate	6.0%
MM (as in Example 1)	1.0%
VM (as in Example 1)	0.5%
fructo-oligosaccharides	3.0%
microalgae oil DHASCO, Martek Biosciences Corporation	1.0%
sodium carbonate	0.2%
vanilla	0.3%
soya lecithin	0.4%
antioxidant vit E	0.01%

Add soya isolate in 50% of water and pass the blend through a colloid mill. Leave the mix for two hours. Add the sucrose, fructo-oligosaccharide and mineral mix in 30% of water and add to the soya isolate blend and heat it up to 65°C in a tank. Add emulsifier and antioxidant to the microalgal oil, mix shortly and add to blend. Add vitamins and vanilla to the blend, homogenize the mix at 170/50 bar and treat under UHT conditions (141°C for 5 sec.). The mix is then cooled down to 20°C, stored in a sterile tank and aseptically filled into brik type packages.

- 12 -

PCT/EP99/02668

#### Claims:

- 1. A nutritional composition for pregnant and lactating women comprising:
  - a non-milk-derived protein source,
  - a dietary fibre and/or fibre-like substance.
  - a source of polyunsaturated fatty acids,
  - minerals and vitamins recommended for pregnant and lactating women, characterized in that the composition is lactose-free and contains less than 2 % of milk or milk-derived products by weight of dry components of the composition.
- 2. The composition according to claim 1, comprising the following minerals and vitamins: calcium, magnesium, iron, copper, zinc, iodine, vitamin A, vitamin E, vitamin B<sub>1</sub>, vitamin B<sub>2</sub>, vitamin B<sub>6</sub>, vitamin B<sub>12</sub>, vitamin C, folic acid, niacin.
- 3. The composition according to claim 1 or claim 2 comprising the following daily amounts of essential nutrients:
  - 1.0 mg (preferably 1.0 g) to 50 g of a non-milk-derived protein source, preferably 8 to 15 g, particularly preferred 10 g,
  - 1.0 mg to 30 g of dietary fibre and/or fibre-like substance, preferably 4 to 6 g, particularly preferred 5 g,
  - of polyunsaturated fatty acids
  - 0.01 μg (preferably 0.1 μg) to 2.0 g of long-chain polyunsaturated fatty acids, preferably 0.1 to 0.4 g, particularly preferred 0.25 g and/or
  - 0.5 g to 12 g of alpha-linolenic acid, preferably 0.5 to 2.0 g, particularly preferred 1.0 g and/ or
  - 1 g to 30 g of linoleic acid, preferably 2.0 to 5.0 g, particularly preferred 3.0 g,
  - 0.1 mg to 3.6 g calcium, preferably 320 to 530 mg, particularly preferred 400 mg,
  - 0.1 mg to 1.1 g magnesium, preferably 60 to 90 mg, particularly preferred 75 mg,
  - 0.01 mg to 90 mg iron, preferably 8 to 12 mg, particularly preferred 10 mg,
  - 0.01 mg to 60 mg zinc, preferably 4 to 6 mg, particularly preferred 5 mg,
  - 0.01 mg to 3 mg copper, preferably 0.1 to 0.4 mg, particularly preferred 0.25 mg,
  - 0.01 mg to 0.6 mg iodine, preferably 24 to 36 μg, particularly preferred 30 μg,
  - 0.01 mg to 4.0 mg vitamin A, preferably 80 to 120 μg, particularly preferred 100 μg,

- 13 -

0.01 mg to 40 mg vitamin E, preferably 2.4 to 3.6 mg, particularly preferred 3 mg, 0.01 mg to 5 mg vitamin  $B_1$ , preferably 0.16 to 0.24 mg, particularly preferred 0.2 mg, 0.01 mg to 6 mg vitamin  $B_2$ , preferably 0.32 to 0.48 mg, particularly preferred 0.4 mg, 0.01 mg to 7 mg vitamin  $B_6$ , preferably 0.24 to 0.36 mg, particularly preferred 0.3 mg, 0.01  $\mu$ g to 8  $\mu$ g vitamin  $B_{12}$ , preferably 0.4  $\mu$ g to 0.6  $\mu$ g, particularly preferred 0.5  $\mu$ g, 0.01 mg to 300 mg vitamin C, preferably 20 to 30 mg, particularly preferred 25 mg, 1.0  $\mu$ g to 1.2 mg folic acid, preferably 160 to 240  $\mu$ g, particularly preferred 200  $\mu$ g, and

0.01 mg to 60 mg niacin, preferably 1.6 to 2.4 mg, particularly preferred 2 mg.

- 4. The composition according to any one of the preceding claims further comprising one or more of the following minerals and vitamins: potassium, selenium, manganese, vitamin K, biotin, pantothenic acid, choline, vitamin D3 and beta-carotene.
- 5. The composition according to any one of the preceding claims wherein the non-milk derived protein source is selected from one or more members of the group consisting of egg protein, soy protein (particularly soy protein low in isoflavones), soy isolate, pea protein, pea hydrolysate, rice protein, rice hydrolysate, soy hydrolysate, wheat hydrolysate, potato hydrolysate, rapeseed protein, lupin protein, wheat protein and wheat isolate.
- 6. The composition according to any one of the preceding claims, wherein the fibre-like is an oligosaccharide, in particular a fructo-oligosaccharide and/or a xylo-oligosaccharide.
- The composition according to any one of the preceding claims wherein the daily amounts of essential nutrients are provided for in a single dose unit.
- 8. A food product or dietary supplement comprising the composition according to any one of the preceding claims.

## INTERNATIONAL SEARCH REPORT

International Application No

_			PCI/EP 99	702668					
A. CLASSI IPC 6	FICATION OF SUBJECT MATTER A23L1/308 A23L1/30 A23L1/30	05							
According to	o international Patent Classification (IPC) or to both national classific	ation and IPC							
B. FIELDS	SEARCHED								
Minimum do IPC 6	ocumentation searched (classification system followed by classificati A23L	on symbols)							
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched									
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)									
C. DOCUME	ENTS CONSIDERED TO BE RELEVANT								
Category *	Citation of document, with indication, where appropriate, of the rel		Relevant to daim No.						
X	US 4 737 364 A (KALOGRIS THEODORE 12 April 1988 (1988-04-12) column 3, line 5 - line 24; exam			1-8					
X	H. SCHERZ, F. SENSER: "Food Compand Nutrition Tables 1989/90" 1989 , WISSENSCHAFTLICHE VERLAGSGESELLSCHAFT MBH , STUTTGA XP002114112 pages 679, 680, 702-705, 708, 709 809	ART		1-8					
		<del> </del>							
X Furth	er documents are listed in the continuation of box C.	X Patent family m	nembers are fisted	n annex.					
•	egories of cited documents : nt defining the general state of the art which is not	"T" later document public or priority date and cited to understand	not in conflict with	the application but					
considered to be of particular relevance Inverse earlier document but published on or after the international "X" docu			ar relevance; the c	almed invention					
ruling date  "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  "C" document of particular relevance; the document is taken alone document of particular relevance; the claimed invention contained to involve an inventive step when the considered to involve an inventive step when the									
other n	re other such docu- is to a person skilled amily								
Date of the a	actual completion of the international search	Date of mailing of th	<del></del>						
2	September 1999	20/09/19	99.						
Name and m	nalling address of the ISA European Patent Office, P.B. 5818 Patentiaan 2 NL – 2280 HV Rijswijk	Authorized officer							
	Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Bendl, E	:						

### INTERNATIONAL SEARCH REPORT

International Application No
PC ./EP 99/02668

		PC./EP 99/02668		
C.(Continu	ation) DOCUMENTS CONSIDERED TO BE RELEVANT			
Category '	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
X	DATABASE WPI Section Ch, Week 8731 Derwent Publications Ltd., London, GB; Class D13, AN 87-219994 XP002114114 & SU 1 274 665 A (ISTRINSK DAIRY IND), 7 December 1986 (1986-12-07) abstract	1-8		
Υ	EP 0 705 539 A (FRIESLAND BRANDS BV) 10 April 1996 (1996-04-10) page 2, line 10 - line 17 page 3, line 14 - line 16	1-8		
Y	DEUTSCHE GESELLSCHAFT FÜR ERNÄHRUNG: "Empfehlungen für die Nährstoffzufuhr" 1991 , UMSCHAU VERLAG , GERMANY XP002114113 pages 146-149	1-8		
Y	DATABASE CAB 'Online! CAB INTERNATIONAL, WALLINGFORD, OXON, GB BUSINCO L: "Prophylaxis of cow's milk allergy." retrieved from DIALOG Database accession no. 00547049 XP002114127 abstract	1-8		

### INTERNATIONAL SEARCH REPORT

nformation on patent family members

International Application No
PC./EP 99/02668

Patent document cited in search repor	t	Publication date	Patent family member(s)	Publication date
US 4737364	Α	12-04-1988	NONE	
SU 1274665	Α	07-12-1986	NONE	
EP 0705539	Α	10-04-1996	NL 9401644 A CN 1125514 A NL 1001372 C NL 1001372 A	01-05-1996 03-07-1996 12-09-1996 11-07-1996

Fo-em PCT/ISA/210 (patent family annex) (July 1992)